



Envisat GDR Cross calibration Report

Cycle 043

28-11-2005 02-01-2006

Prepared by :	Y. Faugere, CLS F. Mertz, CLS J. Dorandeu, CLS	
Accepted by :	J. Dorandeu, CLS	
Approved by :	N. Picot, CNES	



1 Introduction. Document overview

The purpose of this document is to report the major features of the cross-calibration between Envisat and the ERS-2 and Jason-1 missions. The document is associated with data dissemination on a cycle by cycle basis.

The objectives of this document are :

- To present the major useful cross-calibration results for the current cycle

- To report any change likely to impact the comparison between Envisat and other missions, from instrument status to software configuration

It is divided into the following topics:

- Cycle overview**

- Cross Calibration with ERS-2**

- Cross Calibration with Jason-1**

2 Cycle overview

Envisat cycle 043 has been produced with the IPF processing chain V5.02 and the CMA Reference Software V7.1_07. The content of this science software version is described in a document available on the ESA PCS web site ([3]). The Envisat quality assessment report ([4]) summarizes the major features of the Envisat data quality for this cycle of data.

The cross-calibration with ERS-2 OPR2 version 6.5 from CERSAT centre has been performed with ERS-2 OPR cycle 111. The main results for cycle 111 are reported in the ERS-2 Quality assessment report [10]. All the necessary updates were performed on ERS-2 data to be homogeneous with the Envisat data set.

The cross-calibration with Jason-1 GDRs (CMA Reference Software V7.1_05 and V7.1_06) has been performed with Jason-1 GDRs cycles 143 to 147. The content of this science software version is described by N.Picot (electronic communication, October 21, 2005) [12]). The Jason-1 quality assessment report ([1]) summarizes the major features of the Jason-1 data quality for these cycle of data.

3 Cross Calibration with ERS-2

The Envisat/ERS-2 cross-calibration results are not available for this cycle of data.

4 Cross Calibration with Jason-1

Jason-1 GDRs data (cycle 143 to 147) are used for this cross calibration. The parameters used to compute the sea surface height (SSH) for Envisat and Jason-1 are:

- Ku range (ocean retracking)
- POE orbit
- Dual frequency ionospheric correction
- MWR derived wet troposphere correction
- ECMWF dry tropospheric correction
- Non parametric sea state bias
- MOG2D
- Total geocentric GOT00 ocean tide height
- Geocentric pole tide height
- Solid earth tide height

Note that within the IPF version 5.02, the actual value of Ultra Stable Oscillator clock period is used within the L1b processing instead of the nominal one as it was used in previous IPF versions. This evolution implies a +2.5 cm jump on the Envisat SSH between cycle 40 and 41. To avoid this jump, and correct for the USO drift, users are advised to apply the correction provided by ESA on cycles 9 to 40 ([9]).

Some comparisons were also performed using the ECMWF wet troposphere correction for both Envisat and Jason-1, to prevent possible discrepancies from radiometer corrections.

Several analyses were performed for this cross calibration study:


- comparison of altimeter and radiometer parameters
- comparison of Sea Level Anomalies relative to a Mean Sea Surface
- computation of a long wavelength error on Envisat
- comparison on a same time/space sampling

10 day crossovers are used to compare SSH estimations from Envisat and Jason-1 while shorter time lags (3 hours) are selected for altimeter and radiometer parameters.

4.1 Dual-crossover points

4.1.1 3-hour and 10-day crossover points location


For Envisat Cycle 043 the location of crossover points with 3-hour and 10 day time lags between Envisat and Jason-1 are given on the following figures:



Most of the crossover points are located at high latitude. With 3-hour time lag there are only a few crossover points at mid and low latitudes. This geographical pattern is not constant for every Envisat cycle since Jason-1 is not sun-synchronous. When more Envisat data become available, (Jason-1/Envisat) comparisons will be performed over 12 Jason-1 cycle windows, so that the geographical sampling by Jason-1/Envisat crossovers will be constant.

4.1.2 [Envisat - Jason-1] Ku-band SWH differences

The scatter plot of crossover points with 3-hour time lag between Envisat and Jason-1 Ku-band SWH measurements is given on the following figure:




Analysis	Number	Mean (m)	Std. dev. (m)
EN-J1 SWH (m)	1176	0.13	0.25

There is a small bias between the two satellites: Envisat waves are slightly higher than Jason-1 ones.

4.1.3 [Envisat - Jason-1] Ku-band Sigma0 differences

The scatter plot of crossover points with 3-hour time lag between Envisat and Jason-1 Ku-band Sigma0 measurements is given on the following figure:




Analysis	Number	Mean (dB)	Std. dev. (dB)
EN-J1 Sigma0 (dB)	1176	-2.91	0.45

Jason-1 Ku-band sigma0 is 2.8 dB higher than Envisat. Envisat Ku-band sigma0 has been aligned on ERS-2 to satisfy the MWC wind model. Notice that Jason-1 Ku-band sigma0 is 2.3 dB higher than TOPEX. This difference is described in (Vincent et al., 2003 [14]).

4.1.4 [Envisat - Jason-1] radiometer wet troposphere differences

The scatter plot of crossover points with 3-hour time lag between Envisat and Jason-1 radiometer wet troposphere correction is given on the following figure:



Analysis	Number	Mean (cm)	Std. dev. (cm)
EN-J1 radiometer wet troposphere correction (m)	1176	0.78	1.18

Results are consistent over dry areas. There are not enough crossover points at low latitudes to comment the differences in wet areas.

4.1.5 [Envisat - Jason-1] SSH differences


[Envisat - Jason-1] SSH differences at crossover points with 10 day time lag are computed in two configurations:

- using the radiometer wet troposphere correction
- using the ECMWF wet troposphere correction

When using a selection to remove shallow waters (1000 m), global statistics are:

Analysis	Number	Mean (cm)	Std. dev. (cm)
EN-J1 SSH	97627	30.27	6.45
EN-J1 SSH with ECMWF wet troposphere	97627	31.14	6.51


The differences are plotted on the following figure (data are centered about the mean value):




The two maps are very close. There are small scale [Envisat - Jason-1] differences in high variability areas, but also large scale differences in the Pacific ocean.

The cycle by cycle mean and standard deviation of [Envisat-Jason-1] differences of SSH at 10-day dual crossover using the ECMWF wet troposphere correction are plotted in the following figure:

Mean of X_SSH_TRO_HUM_ECMWF cycle




Standard deviation of X_SSH_TRO_HUM_ECMWF per cycle




The mean difference decreases during the first year of Envisat (cycles 10-20). Then, the difference stabilizes around 31 cm on cycle 20 onwards. The standard deviation of the difference is reduced on cycle 41 due to the new ground segment configuration.

4.2 SLA Comparisons

Envisat and Jason-1 Sea Level anomalies relative to CLS01 Mean Sea Surface are computed. Global statistics are computed over deep ocean areas (1000 m) and low variability. In order to see fine features, maps are centered about the mean value.



Analysis	Number	Mean (cm)	Std. dev. (cm)
Envisat SLA	1207285.0000000	49.28	9.52



Analysis	Number	Mean (cm)	Std. dev. (cm)
Jason-1 SLA	1635550.0000000	18.81	9.17


There is a very good correlation between the two maps. The SLA standard deviation for both

Envisat and Jason-1 is about 9.5 cm. Differences are mainly due to the spatial and temporal sampling of the ocean.

4.3 Long wavelength error reduction

4.3.1 Long wavelength error

The Envisat long wavelength error has been computed by global minimization of (EN-J1) SSH differences. The method is described in (Le Traon et al., 1998 [7]). The map of the error is plotted on the following figure (data are centered about the mean value):



Analysis	Number	Mean (cm)	Std. dev. (cm)
Envisat lw error	1364664.0000000	30.53	2.48

The estimated long wavelength error has a small variance which confirms the good quality of the Envisat orbit.


4.3.2 Impact on crossover performances

Global statistics for 35 days [Envisat - Envisat] and 10 days [Envisat - Jason-1] are only computed over deep ocean areas (1000 m) :

Analysis	Number	Mean (cm)	Std. dev. (cm)
EN/EN SSH	37373	-0.71	8.14
EN/EN SSH with orbit error	37373	-0.08	7.73

Analysis	Number	Mean (cm)	Std. dev. (cm)
EN-J1 SSH	97627	30.27	6.45
EN-J1 SSH with orbit error	97627	0.06	6.07


The [Envisat - Jason-1] difference corrected for the estimate Envisat long wavelength error are plotted on the following figure (data are centered about the mean value):



The large scale differences in the Pacific ocean are noticeably reduced.

4.3.3 Impact on SLA performance

Envisat Sea Level anomalies relative to CLS01 Mean Sea Surface using the long wavelength error are computed. Global statistics are computed using a selection to remove shallow waters (1000 m). Map is centered about the mean value.



Analysis	Number	Mean (cm)	Std. dev. (cm)
Envisat SLA	1207285.0000000	18.69	9.49

The slight impact on Envisat SLA variance shows that the Envisat long wavelength error is low.


4.4 Comparison on a same time/space sampling

Envisat and Jason-1 are now compared on a same time/space sampling:

- 35 day period
- $|\text{latitude}| < 66$


4.4.1 Rms of Ku-band range statistics

The histograms of Envisat and Jason-1 Rms of Ku-band range are given on the following figures:
Envisat RMS of Ku-band range, $|\text{Latitude}| < 66$ (unit : cm)



Global nb of points :	1309794	Sel. nb of points :	1309794	Sample interval :	0.200
Global mean :	8.858	Selected mean :	8.858	Maximum value :	25.000
Global Std :	2.717	Selected std :	2.717	Minimum value :	1.900


Jason-1 RMS of Ku-band range (unit : cm)




Global nb of points :	1838778	Sel. nb of points :	1838778	Sample interval :	0.200
Global mean :	7.804	Selected mean :	7.804	Maximum value :	20.000
Global Std :	2.270	Selected std :	2.270	Minimum value :	0.520

4.4.2 Ku-band SWH statistics

The histograms of Envisat and Jason-1 Ku-band SWH are given on the following figures:




Global nb of points	: 1309794	Sel. nb of points	: 1309794	Sample interval	: 0.050
Global mean	: 2.576	Selected mean	: 2.576	Maximum value	: 10.993
Global Std	: 1.239	Selected std	: 1.239	Minimum value	: 0.000




Global nb of points	: 1838778	Sel. nb of points	: 1838778	Sample interval	: 0.050
Global mean	: 2.469	Selected mean	: 2.469	Maximum value	: 10.994
Global Std	: 1.262	Selected std	: 1.262	Minimum value	: 0.000

The cycle per cycle mean of Ku-band SWH measurements for Envisat and Jason-1 is plotted as a function of the cycle number on the following figure:





The cycle per cycle mean difference of Ku-band SWH measurements between Envisat and Jason-1 is plotted as a function of the cycle number on the following figure:




4.4.3 Ku-band Sigma0 statistics

The histograms of Ku-band Sigma0 for Envisat and Jason-1 are given on the following figures:




The general shape of the Envisat histogram is not significantly different from the one obtained at global scale.

The cycle per cycle mean of Ku-band Sigma0 measurements for Envisat and Jason-1 is plotted as a function of the cycle number on the following figure:




The cycle per cycle mean difference of Ku-band Sigma0 measurements between Envisat and Jason-1 is plotted as a function of the cycle number on the following figure:




4.4.4 Troposphere statistics

The histograms of Envisat and Jason-1 radiometer wet troposphere correction are given on the following figures:



Global nb of points :	1309794	Sel. nb of points :	1309794	Sample interval :	0.200
Global mean :	-15.097	Selected mean :	-15.097	Maximum value :	-0.100
Global Std :	9.428	Selected std :	9.428	Minimum value :	-50.000



Global nb of points :	1838778	Sel. nb of points :	1838778	Sample interval :	0.200
Global mean :	-14.641	Selected mean :	-14.641	Maximum value :	-0.100
Global Std :	9.182	Selected std :	9.182	Minimum value :	-49.990

4.4.5 SSH crossover performances

10-day crossover points are computed for both Jason-1 and Envisat. Global statistics of SSH differences at crossovers are computed using a selection to remove shallow waters (1000 m):

Analysis	Number	Mean (cm)	Std. dev. (cm)
EN/EN SSH	16911	-0.62	6.55

Analysis	Number	Mean (cm)	Std. dev. (cm)
J1/J1 SSH	20268	-0.39	6.06


Using an additional selection to remove areas of high ocean variability and high latitudes (> 50 deg) leads to:

Analysis	Number	Mean (cm)	Std. dev. (cm)
EN/EN SSH	10813	-0.43	6.16

Analysis	Number	Mean (cm)	Std. dev. (cm)
J1/J1 SSH	9291	-0.49	5.68

The cycle per cycle standard deviation of SSH measurements is plotted as a function of the cycle number on the following figure:

Std dev. of crossover points, |Latitude|<50°, Bathy<-1000m, Var<20cm



These results show comparable performances for Envisat and Jason-1.

4.4.6 SLA relative to MSS

Envisat and Jason-1 Sea Level anomalies relative to CLS01 Mean Sea Surface are computed. Global statistics are computed removing shallow waters (1000 m) and areas of high ocean variability (20 cm).

Analysis	Number	Mean (cm)	Std. dev. (cm)
Envisat SLA	1207644.000000	49.32	10.08

Analysis	Number	Mean (cm)	Std. dev. (cm)
Jason-1 SLA	1635550.000000	18.81	9.17

These results show comparable performances in terms of SLA variability (standard deviation), and also confirm the crossover estimation of the (Envisat-Jason-1) bias.

References

- [1] Ablain M. et al.: Jason-1 GDR quality assessment report, Cycle 143 to 147. *Technical note ALP-RP-P2-EX-21072-CLS* Available at http://www.aviso.oceanobs.com/html/donnees/calval/validation_report/j1/welcome_uk.html
- [2] Dorandeu J., 2000: Note on ERS-2 Sigma0 variations since January 2000. *Technical note CLS/DOS/NT/00.286*
- [3] EOO/EOX, October 2005, Information to the Users regarding the Envisat RA2/MWR IPF version 5.02 and CMA 7.1 Available at <http://earth.esa.int/pcs/envisat/ra2/articles/>
- [4] Faugere Y. et al.: Envisat GDR quality assesement report (cyclic), Cycle 043, *Technical note SALP-RP-P2-EX-21121-CLS043* Available at http://www.aviso.oceanobs.com/html/donnees/calval/validation_report/en/welcome_uk.html
- [5] Gaspar P. and F. Ogor, 1996: Estimation and analysis of the sea state bias of the new ERS-1 and ERS-2 altimetric data (version6). *Report of task 2 of IFREMER Contract N 96/2.246 002/C.*
- [6] Labroue S. and E. Obligis, 2003: Neural network retrieval algorithms for the ENVISAT/MWR. *Technical note CLS.DOS/NT/03.848*
- [7] Le Traon,P.-Y., F. Ogor, 1998: ERS-1/2 orbit improvement using TOPEX/POSEIDON: The 2cm challenge. *Journal of Geophys. Res., COL. 103, NO. C4, pages 8045-8057*
- [8] Martini A., and P. Fmnia, 2000: The ERS SPTR2000 altimetric range correction: Resultats and validation. *ERE-TN-ADQ-GSO-6001*
- [9] Martini A., 2003: Envisat RA-2 Range instrumental correction : USO clock period variation and associated auxiliary file, Technical Note ENVI-GSEG-EOPG-TN-03-0009 Available at http://earth.esa.int/pcs/envisat/ra2/articles/USO_clock_corr_aux_file.pdf
- [10] Mertz F. et al.: Validation of ERS-2 OPR Cycle 111. *Technical note CLS.OC.NT/03.702 issue 111* Available at <http://www.ifremer.fr/cersat/en/documentation/references/oprmon.htm>
- [11] Obligis E., L. Eymard, N. Tran, 2003: ERS-2/MWR drift evaluation and correction. *Technical note CLS.DOS/NT/03.688*
- [12] Picot N., October 21, 2005: New Jason-1 operational production chain. *Electronic communication.*
- [13] Scharroo R. and P. N. A. M. Visser, 1998: Precise orbit determination and gravity field improvement for the ERS satellites. *J. Geophys. Res., 103, C4, 8113-8127*
- [14] Vincent,P., Desai S.D., Picot N. and Case K., 2003: The first generation of IGDRs and GDRs products to be made available after completion of the Jason-1 verification phase. *Memo to Jason-1 PIs and CoIs.*