



TOPEX/Poseidon MGDR Quality Assessment Report

Cycle 391

26-04-2003 06-05-2003

Prepared by :	C. Schgounn, CLS G. Pontonnier, CLS M. Ablain, CLS	
Accepted by :	J. Dorandeu, CLS	
Quality visa :	M. Destouesse, CLS	
Approved by :	N. Picot, CNES	



CENTRE NATIONAL D'ETUDES SPATIALES



JPL
Jet Propulsion Laboratory
California Institute of Technology

SALP-RP-P2-EX-21120-CLS391

Edition 01.0, July 2003

1 Introduction. Document overview

The purpose of this document is to report the major features of the data quality from the Topex/Poseidon mission. The document is associated with data dissemination on a cycle by cycle basis.

The objectives of this document are :

- To provide a data quality assessment
- To provide users with necessary information for data processing
- To report any change likely to impact data quality at any level, from instrument status to software configuration
- To present the major useful results for the current cycle

It is divided into the following topics:

- [**Cycle overview**](#)
- [**CALVAL main results**](#)

2 Cycle overview

2.1 Cycle quality and performances

Data quality for this cycle appears to be nominal. For this cycle, the crossover standard deviation is 7.02 cm rms, and the standard deviation of Sea Level Anomalies (SLA) relative to a Mean Sea Surface is 9.88 cm.

2.2 Warnings and recommendations

- Tape recorder failures : There is a lot of data gaps due to tape recorder anomalies, especially in the Indian Ocean.
- Problems in the interpolation of the TMR parameters : 3.93% of the measurements are removed by the TMR correction criterion, mainly due to tape recorder failures.


3 CALVAL main results

This section presents results that illustrate data quality during this cycle. These verification products are produced operationally so that they allow systematic monitoring of the main relevant parameters.

3.1 Missing measurements

703084 altimeter measurements are present, and 91480 are missing.

The map below shows all the available measurements for this cycle and illustrates the tape recorder problems. The latter figure shows missing 1Hz measurements in the GDRs, with respect to a 1 Hz sampling of a nominal repeat track.




3.2 M-GDR quality flags


The following table indicates the percentage of measurements for which those flags are set.

Name	Description	% bad
Geo_Bad_1	altimeter land flag	25.78
Geo_Bad_1	ice flag	4.64
Geo_Bad_1	radiometer land flag	27.70
Alt_Bad_1	conditions 1 altimeter	4.55
Alt_Bad_2	conditions 2 altimeter	4.42
Geo_Bad_2	rain (liquid water in excess)	6.57
Geo_Bad_2	less than 4 points for CSR3.0 tide calculation	0.36
Geo_Bad_2	less than 4 points for FES95.2.1 tide calculation	2.49
TOPEX	TOPEX not valid	0.00
TMR	TMR not valid	0.00
TMR_Bad	Brightness temperatures not valid	4.94
DORIS	DORIS not valid	0.00


3.3 M-GDR parameter plots




Cycle 391 – GDR_M – TOPEX – All passes –







Cycle 391 – GDR_M – TOPEX – All passes –




Cycle 391 – GDR_M – TOPEX – All passes –









3.4 Editing

The following table gives for each tested parameter, minimum and maximum thresholds, the number and the percentage of points removed.

As a comparison, the mean percentage over one year (1997) is also given.


There are problems in the interpolation of the TMR parameters since cycle 371 when there are missing measurements (tape recorder failures). These bad measurements are removed by the TMR correction criterion but some of them have been kept. Thus a new criterion has been added to the editing procedure since the cycle 376 to remove all the measurements where the absolute value of the difference between the TMR correction and the ECMWF model wet tropospheric correction is greater than 20 cm.

The number and percentage of points removed by each criterion is given on the following table. Note that these statistics are obtained with measurements already edited for radiometer land flag (27.70 % of points removed) and ice flag (4.64 % of points removed).


Parameters	Min Thres.	Max Thres.	Unit	Mean removed in 1997	% removed
Sea surface height	-130.000	100.000	m	1.37	0.44
Number of 20/10Hz valid points Poseidon/TOPEX	5.000	-		1.37	0.62
Std. deviation of range	0.000	0.100	m	1.85	1.39
Off nadir angle from waveform	0.000	0.400	deg	1.36	3.55
Dry tropospheric correction	-2.500	-1.900	m	0.00	0.00
Invert barometer correction	-2.000	2.000	m	0.00	0.00
TMR wet tropospheric correction	-0.500	-0.001	m	0.34	3.93
Ionospheric correction (Poseidon:Doris, TOPEX:Dual)	-0.400	0.040	m	0.00	0.00
Significant wave height	0.000	11.000	m	1.46	0.27
Sea state Bias	-0.500	0.000	m	1.39	0.41
Backscatter coefficient	7.000	30.000	dB	1.44	0.38
Ocean tide height	-5.000	5.000	m	0.01	0.52
Earth tide	-1.000	1.000	m	0.00	0.00
Pole tide	-15.000	15.000	m	0.00	0.00
TMR and ECMWF tropospheric differences	-0.200	0.200	m	NaN	0.50
Spline fitting					0.01

The following three maps are complementary: they show respectively the removed, the selected measurements and the percentage of selected measurements in the editing procedure.


Valid data
TOPEX/Poseidon Cycle 391 (26/04/2003 / 06/05/2003)



Edited measurements
TOPEX Cycle 391 (26/04/2003 / 06/05/2003)




Percentage of valid data relative to the nominal pass
TOPEX/Poseidon Cycle 391 (26/04/2003 / 06/05/2003)




3.5 Ionospheric correction


TOPEX/Poseidon Cycle 391 (26/04/2003 – 06/05/2003), ascending passes




TOPEX/Poseidon Cycle 391 (26/04/2003 – 06/05/2003), descending passes




TOPEX/Poseidon Cycle 391 (26/04/2003 – 06/05/2003), ascending passes




TOPEX/Poseidon Cycle 391 (26/04/2003 – 06/05/2003), descending passes




TOPEX/Poseidon Cycle 391 (26/04/2003 – 06/05/2003), ascending passes




TOPEX/Poseidon Cycle 391 (26/04/2003 – 06/05/2003), descending passes



3.6 Wet tropospheric corection



3.7 Crossover statistics



T1T1 – CROSSOVER STATISTICS

Without orbit error correction

SSH = Corrected sea surface height

RAPPEL DES SELECTIONS

Type de points de croisement: T1T1

Zone géographique (deg): -90 / 90 , 0 / 360

Seuil sur les écarts d'analyse 0.00 (moy)

30.00 (seuil)

Selection(s) sur les champs :

CL Arc 1 :=INTERP_SPLN

CL Arc 2 :=INTERP_SPLN

Seuil Min +: 0.0000000

Seuil Max : 0.0000000

Selection(s) sur les écarts :

Aucune

RESULTATS STATISTIQUES

Valeur minimale : -29.4500

Valeur maximale : 28.5000

Difference Max – Min: 57.9500

Nombre de points lus: 5180


Nombre de points sélectionnés: 5085

Moyenne : 0.184568

Ecart-type : 7.04191

Moyenne Quadratique : 7.04433

CLS Space Oceanography Division



T1T1 – CROSSOVER STATISTICS

With orbit error correction

SSH = Corrected sea surface height

RAPPEL DES SELECTIONS

Type de points de croisement: T1T1

Zone géographique (deg): -90 / 90 , 0 / 360

Seuil sur les écarts d'analyse 0.00 (moy)

30.00 (seuil)

Selection(s) sur les champs :

CL Arc 1 :=INTERP_SPLN

CL Arc 2 :=INTERP_SPLN

Seuil Min +: 0.0000000

Seuil Max : 0.0000000

Selection(s) sur les écarts :

Aucune

RESULTATS STATISTIQUES

Valeur minimale : -29.2100

Valeur maximale : 28.9600

Difference Max – Min: 58.1700

Nombre de points lus: 5180


Nombre de points sélectionnés: 5088

Moyenne : 0.0561930

Ecart-type : 6.22869

Moyenne Quadratique : 6.22894

CLS Space Oceanography Division



T1T1 – CROSSOVER STATISTICS

SSH, BATHY < -1000 m, VAR_OCE < 20 cm, LAT [-50°, +50]

SSH = Corrected sea surface height before orbit error

RAPPEL DES SELECTIONS

Type de points de croisement: T1T1

Zone géographique (deg): -50 / 50 , 0 / 360

Seuil sur les écarts d'analyse : aucun

Selection(s) sur les champs :

CL Arc 1 :=BATHY

CL Arc 2 :=BATHY

Seuil Min : aucun

Seuil Max : -1000000.00

[...]

Selection(s) sur les écarts :

Aucune

RESULTATS STATISTIQUES

Valeur minimale : -26.3600

Valeur maximale : 32.3000

Difference Max – Min: 58.6600

Nombre de points lus: 2740


Nombre de points sélectionnés: 2499

Moyenne : 0.189628

Ecart-type : 5.73977

Moyenne Quadratique : 5.74290


CLS Space Oceanography Division




3.8 SSH variability

3.8.1 Sea Level Anomaly


TOPEX/Poseidon, cycle 391
Period : 26/04/2003 – 06/05/2003



TOPEX/Poseidon, cycle 391
Period : 26/04/2003 – 06/05/2003




TOPEX/Poseidon, cycle 391
Period : 26/04/2003 – 06/05/2003




3.8.2 Comparison to a precise Mean Sea Surface

The CLS (2001) MSS model is used as a reference to compute SLA. The two following maps respectively show the map of Topex SLA relative to the MSS and differences higher than a 30 cm threshold (after centering the data). The latter figure shows that higher differences are located in high ocean variability areas, as expected.



3.9 Dynamic topography


TOPEX/Poseidon, cycle 391
Period : 26/04/2003 – 06/05/2003



3.10 Wind and wave maps

These two figures show wind and wave estimations derived from 10 days of altimeter measurements.

TOPEX/Poseidon, cycle 391
Period : 26/04/2003 – 06/05/2003



TOPEX/Poseidon, cycle 391
Period : 26/04/2003 – 06/05/2003

