



TOPEX/Poseidon MGDR Quality Assessment Report

Cycle 404

02-09-2003 12-09-2003

Prepared by :	C. Schgounn, CLS G. Pontonnier, CLS M. Ablain, CLS	
Accepted by :	J. Dorandeu, CLS	
Quality visa :	M. Destouesse, CLS	
Approved by :	N. Picot, CNES	



CENTRE NATIONAL D'ETUDES SPATIALES



JPL
Jet Propulsion Laboratory
California Institute of Technology

SALP-RP-P2-EX-21120-CLS404

Edition 01.0, December 2003

1 Introduction. Document overview

The purpose of this document is to report the major features of the data quality from the Topex/Poseidon mission. The document is associated with data dissemination on a cycle by cycle basis.

The objectives of this document are :

- To provide a data quality assessment
- To provide users with necessary information for data processing
- To report any change likely to impact data quality at any level, from instrument status to software configuration
- To present the major useful results for the current cycle

It is divided into the following topics:

- [**Cycle overview**](#)
- [**CALVAL main results**](#)

2 Cycle overview

2.1 Cycle quality and performances

Data quality for this cycle appears to be nominal. For this cycle, the crossover standard deviation is 6.30 cm rms, and the standard deviation of Sea Level Anomalies (SLA) relative to a Mean Sea Surface is 9.34 cm.

2.2 Warnings and recommendations

- Missing measurements :
 - There is a lot of data gaps due to tape recorder anomalies, especially in the Indian Ocean, in the South Pacific Ocean close to the South and Central America coasts and below the Groenland coasts.
 - Pass 179 is missing from current data set. No information allows to explain this anomaly.
- Doris ionospheric correction :


The DORIS instrument experienced an incident on 06,September 2003. Therefore there is no DORIS data available from 05/09/2003(pass 74) to 10/09/2003(pass 227). Only bent ionospheric correction is available during this period.
- Measurements edited by the TMR parameters :

The following anomalies are explained by the problems in the interpolation of the TMR parameters due to tape recorder failures :

 - 2.27% of the measurements are removed by the TMR correction criterion (see the following figure).
 - Some measurements have radiometer land flag set to valid over land. A new criterion has been added to the editing procedure to remove all these measurements (see [Editing](#)) .

The pass 142 is partly rejected (in the southern Indian Ocean) by the radiometer land flag criterium. This flag is set to defalut value. This anomaly seems not linked with the problems in the interpolation of the TMR parameters and is under investigation.

Edited parameter : Radiometer wet tropospheric correction
T/P Cycle 404 (02/09/2003 / 12/09/2003)




3 CALVAL main results

This section presents results that illustrate data quality during this cycle. These verification products are produced operationally so that they allow systematic monitoring of the main relevant parameters.

3.1 Missing measurements

694964 altimeter measurements are present, and 99615 are missing.

The map below shows all the available measurements for this cycle and illustrates the tape recorder problems. The latter figure shows missing 1Hz measurements in the GDRs, with respect to a 1 Hz sampling of a nominal repeat track.





3.2 M-GDR quality flags

The following table indicates the percentage of measurements for which those flags are set.


Name	Description	% bad
Geo_Bad_1	altimeter land flag	25.54
Geo_Bad_1	ice flag	8.07
Geo_Bad_1	radiometer land flag	27.68
Alt_Bad_1	conditions 1 altimeter	5.37
Alt_Bad_2	conditions 2 altimeter	5.20
Geo_Bad_2	rain (liquid water in excess)	5.19
Geo_Bad_2	less than 4 points for CSR3.0 tide calculation	0.45
Geo_Bad_2	less than 4 points for FES95.2.1 tide calculation	3.11
TOPEX	TOPEX not valid	0.00
TMR	TMR not valid	0.00
TMR_Bad	Brightness temperatures not valid	5.25
DORIS	DORIS not valid	0.00

3.3 M-GDR parameter plots










Cycle 404 – GDR_M – TOPEX – All passes –




Cycle 404 – GDR_M – TOPEX – All passes –









3.4 Editing

The following table gives for each tested parameter, minimum and maximum thresholds, the number and the percentage of points removed. As a comparison, the mean percentage over one year (1997) is also given.

There are problems in the interpolation of the TMR parameters since cycle 371 when there are missing measurements (tape recorder failures). These bad measurements are removed by the TMR correction criterion but some of them have been kept. Thus a new criterion has been added to the editing procedure since the cycle 376 to remove all the measurements where the absolute value of the difference between the TMR correction and the ECMWF model wet tropospheric correction is greater than 20 cm.


Probably due to the interpolation problem with the TMR, some measurements have radiometer land flag unset over land. This has no impact on the valid data because these measurements have been edited by the altimetric parameter criteria. Nevertheless, this anomaly leads to wrong statistics of the edited measurements. Therefore a new criterion has been added in the editing procedure to remove all the measurements for which the radiometer land flag is set to ocean and the altimeter land flag is set to land.

The number and percentage of points removed by each criterion is given on the following table. Note that these statistics are obtained with measurements already edited for radiometer land flag (27.68 % of points removed) and ice flag (8.07 % of points removed).


Parameters	Min Thres.	Max Thres.	Unit	Mean removed in 1997	% removed
Sea surface height	-130.000	100.000	m	1.37	0.18
Number of 20/10Hz valid points Poseidon/TOPEX	5.000	-		1.37	0.27
Std. deviation of range	0.000	0.100	m	1.85	1.13
Off nadir angle from waveform	0.000	0.400	deg	1.36	3.56
Dry tropospheric correction	-2.500	-1.900	m	0.00	0.00
Invert barometer correction	-2.000	2.000	m	0.00	0.00
TMR wet tropospheric correction	-0.500	-0.001	m	0.34	2.27
Ionospheric correction (Poseidon:Doris, TOPEX:Dual)	-0.400	0.040	m	0.00	0.00
Significant wave height	0.000	11.000	m	1.46	0.19
Sea state Bias	-0.500	0.000	m	1.39	0.29
Backscatter coefficient	7.000	30.000	dB	1.44	0.26
Ocean tide height	-5.000	5.000	m	0.01	0.15
Earth tide	-1.000	1.000	m	0.00	0.00
Pole tide	-15.000	15.000	m	0.00	0.00
TMR and ECMWF tropospheric differences	-0.200	0.200	m	NaN	0.36
Spline fitting					0.01

The following three maps are complementary: they show respectively the removed, the selected measurements and the percentage of selected measurements in the editing procedure.


Valid data
TOPEX/Poseidon Cycle 404 (02/09/2003 / 12/09/2003)



Edited measurements
TOPEX Cycle 404 (02/09/2003 / 12/09/2003)




Percentage of valid data relative to the nominal pass
TOPEX/Poseidon Cycle 404 (02/09/2003 / 12/09/2003)




3.5 Ionospheric correction


TOPEX/Poseidon Cycle 404 (02/09/2003 – 12/09/2003), ascending passes




TOPEX/Poseidon Cycle 404 (02/09/2003 – 12/09/2003), descending passes




TOPEX/Poseidon Cycle 404 (02/09/2003 – 12/09/2003), ascending passes




TOPEX/Poseidon Cycle 404 (02/09/2003 – 12/09/2003), descending passes




TOPEX/Poseidon Cycle 404 (02/09/2003 – 12/09/2003), ascending passes




TOPEX/Poseidon Cycle 404 (02/09/2003 – 12/09/2003), descending passes



3.6 Wet tropospheric corection



3.7 Crossover statistics



T1T1 – CROSSOVER STATISTICS

Without orbit error correction

SSH = Corrected sea surface height

RAPPEL DES SELECTIONS

Type de points de croisement: T1T1

Zone géographique (deg): -90 / 90 , 0 / 360

Seuil sur les écarts d'analyse 0.00 (moy)
30.00 (seuil)

Selection(s) sur les champs :

CL Arc 1 :=INTERP_SPLN
CL Arc 2 :=INTERP_SPLN
Seuil Min +: 0.0000000
Seuil Max : 0.0000000

Selection(s) sur les écarts :

Aucune

RESULTATS STATISTIQUES

Valeur minimale : -29.9000

Valeur maximale : 29.0100

Difference Max – Min: 58.9100

Nombre de points lus: 4835


Nombre de points sélectionnés: 4726

Moyenne : 0.716007

Ecart-type : 6.31563

Moyenne Quadratique : 6.35608

CLS Space Oceanography Division



T1T1 – CROSSOVER STATISTICS

With orbit error correction

SSH = Corrected sea surface height

RAPPEL DES SELECTIONS

Type de points de croisement: T1T1

Zone géographique (deg): -90 / 90 , 0 / 360

Seuil sur les écarts d'analyse 0.00 (moy)
30.00 (seuil)

Selection(s) sur les champs :

CL Arc 1 :=INTERP_SPLN
CL Arc 2 :=INTERP_SPLN
Seuil Min +: 0.0000000
Seuil Max : 0.0000000

Selection(s) sur les écarts :

Aucune

RESULTATS STATISTIQUES

Valeur minimale : -29.9100

Valeur maximale : 29.8100

Difference Max – Min: 59.7200

Nombre de points lus: 4835


Nombre de points sélectionnés: 4716

Moyenne : 0.223338

Ecart-type : 5.68732

Moyenne Quadratique : 5.69170

CLS Space Oceanography Division



T1T1 – CROSSOVER STATISTICS

SSH, BATHY < -1000 m, VAR_OCE < 20 cm, LAT [-50°, +50]

SSH = Corrected sea surface height before orbit error

RAPPEL DES SELECTIONS

Type de points de croisement: T1T1
 Zone géographique (deg): -50 / 50 , 0 / 360
 Seuil sur les écarts d'analyse : aucun

Selection(s) sur les champs :

CL Arc 1 :=BATHY
 CL Arc 2 :=BATHY
 Seuil Min : aucun
 Seuil Max : -100000.00
 [...]


Selection(s) sur les écarts :

Aucune

RESULTATS STATISTIQUES

Valeur minimale : -31.9700
 Valeur maximale : 30.6700
 Différence Max – Min: 62.6400
 Nombre de points lus: 2672
 Nombre de points sélectionnés: 2448
 Moyenne : 0.365012
 Ecart-type : 5.49846
 Moyenne Quadratique : 5.51056


CLS Space Oceanography Division




3.8 SSH variability

3.8.1 Sea Level Anomaly


TOPEX/Poseidon, cycle 404
Period : 02/09/2003 – 12/09/2003



TOPEX/Poseidon, cycle 404
Period : 02/09/2003 – 12/09/2003




TOPEX/Poseidon, cycle 404
Period : 02/09/2003 – 12/09/2003




3.8.2 Comparison to a precise Mean Sea Surface

The CLS (2001) MSS model is used as a reference to compute SLA. The two following maps respectively show the map of Topex SLA relative to the MSS and differences higher than a 30 cm threshold (after centering the data). The latter figure shows that higher differences are located in high ocean variability areas, as expected.



3.9 Dynamic topography


TOPEX/Poseidon, cycle 404
Period : 02/09/2003 – 12/09/2003



3.10 Wind and wave maps

These two figures show wind and wave estimations derived from 10 days of altimeter measurements.

TOPEX/Poseidon, cycle 404
Period : 02/09/2003 – 12/09/2003



TOPEX/Poseidon, cycle 404
Period : 02/09/2003 – 12/09/2003

