



TOPEX/Poseidon MGRD Quality Assessment Report

Cycle 457

08-02-2005 / 18-02-2005

| | | |
|----------------|--|--|
| Prepared by : | C. Schgounn, CLS G. Pontonnier, CLS M. Ablain, CLS | |
| Accepted by : | J. Dorandeu, CLS | |
| Quality visa : | M. Destouesse, CLS | |
| Approved by : | N. Picot, CNES | |



SALP-RP-P2-EX-21120-CLS457

Edition 01.0, April 2005

1 Introduction. Document overview

The purpose of this document is to report the major features of the data quality from the Topex/Poseidon mission. The document is associated with data dissemination on a cycle by cycle basis.

The objectives of this document are :

- To provide a data quality assessment
- To provide users with necessary information for data processing
- To report any change likely to impact data quality at any level, from instrument status to software configuration
- To present the major useful results for the current cycle

It is divided into the following topics:

[Cycle overview](#)

[CALVAL main results](#)

2 Cycle overview

2.1 Cycle quality and performances

Data quality for this cycle is nominal.

For this cycle, the crossover standard deviation is 6.73 cm rms. When using a selection to remove shallow waters (1000 m), areas of high ocean variability and high latitudes ($> |50|$ deg.) it decreases down to 5.99 cm rms.

The standard deviation of Sea Level Anomalies (SLA) relative to a 7-year Mean Sea Surface is 10.80 cm. When using a selection to remove shallow waters (1000 m), areas of high ocean variability and high latitudes ($> |50|$ deg), it lowers to 9.59 cm .

2.2 Missing measurements

This cycle has no missing pass.

2.3 Warnings and recommendations

- Missing measurements :
 - Since October 08th all mission data recovery requirements have been met via TDRSS real time contacts.
Therefore there is a lot of data gaps, especially in the Indian Ocean, between the East and Southeast Pacific basin, in the South Pacific Ocean close to the South and Central America coasts and below the Groenland coasts.
- Doris switch off :
The DORIS instrument was switched off since the incident on 01, November 2004. All the POE requirements are now met using lasernet tracking data. Only bent ionospheric correction is available.
- Measurements edited by the TMR parameters :
The following anomalies are explained by the problems in the interpolation of the TMR parameters due to tape recorder failures :
 - All measurements of pass 90 are removed by the radiometer land flag criterium.
 - 7.84% of the measurements are removed by the TMR correction criterion.
 - Some measurements have radiometer earth flag set to valid over earth. A new criterion has been added to the editing procedure to remove all these measurements (see [Editing](#)) .

3 CALVAL main results


This section presents results that illustrate data quality during this cycle. These verification products are produced operationally so that they allow systematic monitoring of the main relevant parameters.

3.1 Missing measurements


635300 altimeter measurements are present, and 159348 are missing.

The map below shows all the available measurements for this cycle and illustrates the tape recorder problems. The latter figure shows missing 1Hz measurements in the GDRs, with respect to a 1 Hz sampling of a nominal repeat track.

Available measurements
TOPEX Cycle 457 (08/02/2005 / 18/02/2005)



Missing measurements
TOPEX/Poseidon Cycle 457 (08/02/2005 / 18/02/2005)




3.2 M-GDR quality flags

The following table indicates the percentage of measurements for which those flags are set.


| Name | Description | % bad |
|-----------|---|-------|
| Geo_Bad_1 | altimeter land flag | 25.42 |
| Geo_Bad_1 | ice flag | 2.89 |
| Geo_Bad_1 | radiometer land flag | 27.12 |
| Alt_Bad_1 | conditions 1 altimeter | 4.10 |
| Alt_Bad_2 | conditions 2 altimeter | 3.98 |
| Geo_Bad_2 | rain (liquid water in excess) | 11.07 |
| Geo_Bad_2 | less than 4 points for CSR3.0 tide calculation | 0.37 |
| Geo_Bad_2 | less than 4 points for FES95.2.1 tide calculation | 2.51 |
| TOPEX | TOPEX not valid | 0.00 |
| TMR | TMR not valid | 0.00 |
| TMR_Bad | Brightness temperatures not valid | 11.07 |
| DORIS | DORIS not valid | 0.00 |

3.3 M-GDR parameter plots


Cycle 457 – GDR_M – TOPEX – All passes –




Cycle 457 – GDR_M – TOPEX – All passes –









Cycle 457 – GDR_M – TOPEX – All passes –




Cycle 457 – GDR_M – TOPEX – All passes –









3.4 Editing

The following table gives for each tested parameter, minimum and maximum thresholds, the number and the percentage of points removed. As a comparison, the mean percentage over one year (1997) is also given.

There are problems in the interpolation of the TMR parameters since cycle 371 when there are missing measurements (tape recorder failures). These bad measurements are removed by the TMR correction criterion but some of them have been kept. Thus a new criterion has been added to the editing procedure since the cycle 376 to remove all the measurements where the absolute value of the difference between the TMR correction and the ECMWF model wet tropospheric correction is greater than 20 cm.


Probably due to the interpolation problem with the TMR, some measurements have radiometer land flag unset over land. This has no impact on the valid data because these measurements have been edited by the altimetric parameter criteria. Nevertheless, this anomaly leads to wrong statistics of the edited measurements. Therefore a new criterion has been added in the editing procedure to remove all the measurements for which the radiometer land flag is set to ocean and the altimeter land flag is set to land.

The number and percentage of points removed by each criterion is given on the following table. Note that these statistics are obtained with measurements already edited for radiometer land flag (27.12 % of points removed) and ice flag (2.89 % of points removed).


| Parameters | Min Thres. | Max Thres. | Unit | Mean % removed in 1997 | % removed |
|---|------------|------------|------|------------------------|-----------|
| Sea surface height | -130.000 | 100.000 | m | 1.37 | 0.13 |
| Number of 20/10Hz valid points Poseidon/TOPEX | 5.000 | - | | 1.37 | 0.22 |
| Std. deviation of range | 0.000 | 0.100 | m | 1.85 | 0.85 |
| Off nadir angle from waveform | 0.000 | 0.400 | deg | 1.36 | 2.91 |
| Dry tropospheric correction | -2.500 | -1.900 | m | 0.00 | 0.00 |
| Invert barometer correction | -2.000 | 2.000 | m | 0.00 | 0.00 |
| TMR wet tropospheric correction | -0.500 | -0.001 | m | 0.34 | 7.84 |
| Ionospheric correction (Poseidon:Doris, TOPEX:Dual) | -0.400 | 0.040 | m | 0.00 | 0.23 |
| Significant wave height | 0.000 | 11.000 | m | 1.46 | 0.12 |
| Sea state Bias | -0.500 | 0.000 | m | 1.39 | 0.15 |
| Backscatter coefficient | 7.000 | 30.000 | dB | 1.44 | 0.16 |
| Ocean tide height | -5.000 | 5.000 | m | 0.01 | 0.12 |
| Earth tide | -1.000 | 1.000 | m | 0.00 | 0.00 |
| Pole tide | -15.000 | 15.000 | m | 0.00 | 0.00 |
| TMR and ECMWF tropospheric differences | -0.200 | 0.200 | m | NaN | 0.37 |
| Spline fitting | | | | | 0.01 |

The following three maps are complementary: they show respectively the removed, the selected measurements and the percentage of selected measurements in the editing procedure.


Valid data
TOPEX/Poseidon Cycle 457 (08/02/2005 / 18/02/2005)



Edited measurements
TOPEX Cycle 457 (08/02/2005 / 18/02/2005)




Percentage of valid data relative to the nominal pass
TOPEX/Poseidon Cycle 457 (08/02/2005 / 18/02/2005)




3.5 Ionospheric correction


TOPEX/Poseidon Cycle 457 (08/02/2005 – 18/02/2005), ascending passes



TOPEX/Poseidon Cycle 457 (08/02/2005 – 18/02/2005), descending passes



minimum number of samples per box 1
 maximum number of samples per box 1066



Statistics Y-X

mean = -0.01365
 rms = 0.02057
 std = 0.01539


Order 1 fit polynom

$y = a x + b$
 $a = 0.64167798$
 $b = 0.00803378$


Legend

— Order 1 fit polynom
 — Bisectrix

3.6 Wet tropospheric correction



3.7 Crossover statistics



T1T1 – CROSSOVER STATISTICS

Without orbit error correction

SSH = Corrected sea surface height

RAPPEL DES SELECTIONS


Type de points de croisement: T1T1
 Zone géographique (deg): -90 / 90 , 0 / 360
 Seuil sur les écarts d'analyse DV (moy) 30.00 (seuil)
 Selection(s) sur les champs :
 CL Arc 1 : =INTERP_SPLN
 CL Arc 2 : =INTERP_SPLN
 Seuil Min + : 0.0000000
 Seuil Max : 0.0000000

Selection(s) sur les écarts :
 Aucune

RESULTATS STATISTIQUES

Valeur minimale : -29.4100
 Valeur maximale : 29.8800
 Différence Max – Min: 59.2900
 Nombre de points lus: 3919
 Nombre de points sélectionnés: 3826
 Moyenne : 0.129846
 Écart-type : 6.47580
 Moyenne Quadratique : 6.47711

CLS Space Oceanography Division



T1T1 – CROSSOVER STATISTICS

With orbit error correction

SSH = Corrected sea surface height

RAPPEL DES SELECTIONS


Type de points de croisement: T1T1
 Zone géographique (deg): -90 / 90 , 0 / 360
 Seuil sur les écarts d'analyse DV (moy) 30.00 (seuil)
 Selection(s) sur les champs :
 CL Arc 1 : =INTERP_SPLN
 CL Arc 2 : =INTERP_SPLN
 Seuil Min + : 0.0000000
 Seuil Max : 0.0000000

Selection(s) sur les écarts :
 Aucune

RESULTATS STATISTIQUES

Valeur minimale : -28.0600
 Valeur maximale : 29.3100
 Différence Max – Min: 57.3700
 Nombre de points lus: 3919
 Nombre de points sélectionnés: 3813
 Moyenne : 0.0365041
 Écart-type : 5.48500
 Moyenne Quadratique : 5.48512

CLS Space Oceanography Division



T1T1 – CROSSOVER STATISTICS
SSH, BATHY < -1000 m, VAR_OCE < 20 cm, LAT [-50°, +50]
SSH = Corrected sea surface height before orbit error


RAPPEL DES SELECTIONS

Type de points de croisement: T1T1
 Zone géographique (deg): -50 / 50 , 0 / 360
 Seuil sur les écarts d'analyse : aucun
 Selection(s) sur les champs :
 CL Arc 1 : =BATHY
 CL Arc 2 : =BATHY
 Seuil Min : aucun
 Seuil Max : -100000.00
 CL Arc 1 : =VAR_OCE
 CL Arc 2 : =VAR_OCE
 Seuil Min : aucun
 Seuil Max : 20.000000
 [...]
 Selection(s) sur les écarts :
 Aucune

RESULTATS STATISTIQUES

Valeur minimale : -60.1300
 Valeur maximale : 29.2400
 Différence Max – Min: 89.3700
 Nombre de points lus: 2845
 Nombre de points sélectionnés: 2135
 Moyenne : -0.524070
 Écart-type : 5.99144
 Moyenne Quadratique : 6.01431


CLS Space Oceanography Division




3.8 SSH variability

3.8.1 Sea Level Anomaly


TOPEX/Poseidon, cycle 457
Period : 08/02/2005 – 18/02/2005



TOPEX/Poseidon, cycle 457
Period : 08/02/2005 – 18/02/2005




TOPEX/Poseidon, cycle 457
Period : 08/02/2005 – 18/02/2005




3.8.2 Comparison to a precise Mean Sea Surface

The CLS (2001) MSS model is used as a reference to compute SLA. The two following maps respectively show the map of Topex SLA relative to the MSS and differences higher than a 30 cm threshold (after centering the data). The latter figure shows that higher differences are located in high ocean variability areas, as expected.

TOPEX/Poseidon, cycle 457
Period : 08/02/2005 – 18/02/2005




(SSH - MSS) differences greater than 0.3 m
TOPEX/Poseidon Cycle 457 (08/02/2005 / 18/02/2005)



3.9 Dynamic topography


TOPEX/Poseidon, cycle 457
Period : 08/02/2005 – 18/02/2005



3.10 Wind and wave maps

These two figures show wind and wave estimations derived from 10 days of altimeter measurements.

TOPEX/Poseidon, cycle 457
Period : 08/02/2005 – 18/02/2005



TOPEX/Poseidon, cycle 457
Period : 08/02/2005 – 18/02/2005

